

AN EXTENSION OF ALEXANDROV'S THEOREM ON SECOND DERIVATIVES OF CONVEX FUNCTIONS

J.H.G. FU

ABSTRACT. If f is a function of n variables that is locally uniformly approximable by a sequence of smooth functions satisfying local L^1 bounds on the determinants of the minors of the Hessian, then f admits a second order Taylor expansion almost everywhere. This extends a classical theorem of A.D. Alexandrov, covering the special case in which f is locally convex.

1. INTRODUCTION

A.D. Alexandrov [2] proved that if $f : U \rightarrow \mathbb{R}$ is a locally convex function on a domain $U \subset \mathbb{R}^n$ then f “admits a second derivative almost everywhere”, in the sense that for a.e. $x \in U$ there is a second order Taylor expansion for f at x . In the present article we extend this conclusion to a much larger class of functions. Let us say that f is **twice differentiable at x** if there is a quadratic polynomial $Q_x : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

$$(1) \quad \lim_{y \rightarrow x} \frac{f(y) - Q_x(y)}{|y - x|^2} = 0.$$

Theorem 1.1. *Let $U \subset \mathbb{R}^n$ be open. Suppose $f : U \rightarrow \mathbb{R}$ may be expressed as the locally uniform limit of a sequence $f_1, f_2, \dots$ of smooth functions such that the absolute integrals of all minors of the Hessians of the f_k are locally bounded, i.e.*

$$(2) \quad \int_K \left| \det \left(\frac{\partial^2 f_k}{\partial x_i \partial x_j} \right)_{i \in I, j \in J} \right| \leq C(K), \quad k = 1, 2, \dots$$

whenever $K \subset\subset U$ and $I, J \subset \{1, \dots, n\}$ have the same cardinality. Then f is twice differentiable at a.e. $x \in U$.

(If $n = 1$ then f is approximable in this way iff it is expressible as the difference of two convex functions; but this is not the case for $n \geq 2$, as we show below.)

Alexandrov's theorem is a special case: if f is convex then such a sequence f_k may be obtained by convolution with an approximate identity. But Theorem 1.1 is significantly stronger: even though a convex function f may fail to have first derivatives in the usual sense on a dense set, it has been observed (cf. [1]) that such f admits a multiple-valued differential everywhere, whose graph transforms into the graph of a Lipschitz function under the canonical linear change of variable $(x, y) \mapsto (x + y, x - y)$ of $T^*\mathbb{R}^n \simeq \mathbb{R}^n \times \mathbb{R}^n$. From this point of view, Alexandrov's theorem appears as a consequence of Rademacher's theorem on the almost everywhere differentiability of Lipschitz functions. On the other hand, there exist functions f satisfying the hypothesis of Theorem 1.1 such that the graph of the

Date: August 17, 2009.

differential of f is dense in $T^*\mathbb{R}^n$. In fact, such functions include the Sobolev space $W_{\text{loc}}^{2,n}(U)$ of functions with second distributional derivatives in $L_{\text{loc}}^n(U)$: convolution with an approximate identity yields a sequence of smooth functions with bounded local $W^{2,n}$ norms converging to f in $W_{\text{loc}}^{2,n}$. By Hölder's inequality, the local bounds imply that the resulting sequence satisfies the bounds (2), and the Sobolev embedding theorem implies that the convergence to f is locally uniform. Meanwhile, Hutchinson-Meier [9] observed that if $n \geq 2$ then

$$(3) \quad f_{HM}(x) := x_1 \sin \log \log |x|^{-1}$$

belongs to $W^{2,n}(U)$ for sufficiently small neighborhoods U of 0, while ∇f_{HM} oscillates infinitely often between $(\pm 1, 0, \dots, 0) + o(x)$ as $x \rightarrow 0$ along the x_1 axis. This implies first of all that f_{HM} cannot be expressed as a difference of two convex functions. Furthermore, by cutting off, translating and multiplying by suitable constants, it is easy to construct a $W^{2,n}(\mathbb{R}^n)$ -convergent sum whose differential has a graph that is dense in $T^*\mathbb{R}^n$.

As a final introductory remark, we note that this class of functions is a subset of the class of ‘‘Monge-Ampère functions’’ introduced by Jerrard in [10, 11] (extending the class defined in [5]). It is natural to conjecture that Theorem 1.1 applies to this nominally larger class as well. However, this is just one of many perplexing questions about Monge-Ampère functions: for example, we do not even know that such functions are necessarily continuous, or even locally bounded. On the other hand, it seems highly plausible that every Monge-Ampère function satisfies the hypotheses of Theorem 1.1.

Acknowledgement. I would like to thank Bob Jerrard for stimulating conversations on the topics discussed here.

2. BASIC FACTS

2.1. Some measure theory. Recall that if μ is a Radon measure on $U \subset \mathbb{R}^n$ then its *upper density* at $x \in U$ is

$$\Theta^*(\mu, x) := \limsup_{r \downarrow 0} \frac{\mu(B(x, r))}{\omega_n r^n},$$

and its *density* is

$$\Theta(\mu, x) := \lim_{r \downarrow 0} \frac{\mu(B(x, r))}{\omega_n r^n},$$

if the limit exists. In fact the limit exists for a.e. $x \in U$ with respect to Lebesgue measure, and defines a Lebesgue-integrable function of x , whose integrals yield the absolutely continuous part of μ with respect to the Lebesgue decomposition into absolutely continuous and singular parts (cf. [4], Thm. 3.22).

2.2. Absolute Hessian determinant measures. From now on we fix an open set $U \subset \mathbb{R}^n$, a function $f : U \rightarrow \mathbb{R}$, and an approximating sequence $f_1, f_2, \dots \rightarrow f$ as in the hypothesis of Theorem 1.1. In particular, f is itself continuous. For $d = 0, \dots, n$, and $k = 1, 2, \dots$, we define the measures $\nu_{d,k}$ on U by

$$(4) \quad \nu_{d,k}(S) := \sum_{I, J \subset \{1, \dots, n\}, |I|=|J|=d} \int_S \left| \det \left[\frac{\partial^2 f_k}{\partial x^i \partial x^j} \right]_{i \in I, j \in J} \right|.$$

Taking subsequences, we may assume that the each sequence $\nu_{d,k}, k = 1, 2, \dots$, converges weakly to a Radon measure $\nu_d, d = 0, \dots, n$. We will refer to the ν_d as **absolute Hessian determinant measures** for f .

The following lemma is obvious. For quadratic polynomials $Q : \mathbb{R}^n \rightarrow \mathbb{R}$, put $\|Q\|$ to be the maximum of the absolute values of the coefficients.

Lemma 2.1. *Let Q be a quadratic polynomial. Then $f + Q$ and the approximating sequence $f_1 + Q, f_2 + Q, \dots$ also satisfy the hypothesis of Theorem 1.1. Furthermore the resulting absolute Hessian determinant measures $\tilde{\nu}_d$ for $f + Q$ satisfy*

$$(5) \quad \tilde{\nu}_d \leq d! 8^n \sum_{i+j=d} \|Q\|^i \nu_j, \quad d = 0, \dots, n.$$

2.3. An inequality from multivariable calculus. The key fact that makes the main theorem work is the extension of the following elementary classical inequality about C^2 functions to our larger class. Put ω_n for the volume of the unit ball in $\mathbb{R}^n$.

Lemma 2.2. *If $V \subset\subset U$ is open then*

$$(6) \quad \nu_n(V) \geq \omega_n \left(\frac{\sup_V |f| - \sup_{\partial V} |f|}{\text{diam } V} \right)^n$$

Proof. By the triangle inequality, this is a weaker form of the corresponding statement with f replacing $|f|$, which in turn follows easily from the corresponding statement in the smooth case (cf. [7], Lemma 9.2). To emphasize the point we give a complete proof.

Let $\epsilon > 0$ be given, choose $W \subset\subset V$ so that $\sup_W f - \sup_{\partial W} f > \sup_V f - \sup_{\partial V} f - \epsilon$. Let k_0 be a natural number large enough that $|f(x) - f_k(x)| < \frac{\epsilon}{2}$ for $x \in V$ and $k \geq k_0$. Then for any linear function $\lambda : \mathbb{R}^n \rightarrow \mathbb{R}$ and any $k \geq k_0$

$$(7) \quad \begin{aligned} \sup_W (f_k - \lambda) - \sup_{\partial W} (f_k - \lambda) &\geq \sup_W f_k - \sup_{\partial W} f_k - |\lambda| \text{diam } V \\ &\geq \sup_V f - \sup_{\partial V} f - |\lambda| \text{diam } V - 2\epsilon \end{aligned}$$

Thus if $|\lambda| < \frac{\sup_V f - \sup_{\partial V} f - 2\epsilon}{\text{diam } V}$ then the left hand side is positive, and $f_k - \lambda$ has interior local maximum in W , at which $df_k = \lambda$. Thus for $k \geq k_0$ the image of W under df_k includes the ball of radius $\frac{\sup_V f - \sup_{\partial V} f - 2\epsilon}{\text{diam } V}$, and by the area formula and the properties of weak convergence of measures,

$$\begin{aligned} \nu_n(V) \geq \nu_n(\bar{W}) &\geq \limsup_{k \rightarrow \infty} \int_W |\det D^2 f_k| \\ &\geq \omega_n \left(\frac{\sup_V f - \sup_{\partial V} f - 2\epsilon}{\text{diam } V} \right)^n \end{aligned}$$

by (7), and the lemma follows letting $\epsilon \downarrow 0$. □

In fact we will need to apply Lemma 2.2 to affine subspaces as well.

Corollary 2.3. *Given $d < n$, let $M \subset \mathbb{R}^n$ be a linear subspace of codimension d . For each $x \in M$, denote by P_x the affine d -plane orthogonal to M and passing through the point x . If $V \subset\subset U$ is open then*

$$(8) \quad \nu_d(V) \geq \omega_d \int_M \left| \frac{\sup_{V \cap P_x} |f| - \sup_{\partial(V \cap P_x)} |f|}{\text{diam}(V \cap P_x)} \right|^d dx$$

where the integrand is understood to be zero if $V \cap P_x = \emptyset$.

Proof. Let $W \subset \subset V$. By Lemma 2.2, for each approximating function f_k

$$\begin{aligned} \omega_d \int_M \left| \frac{\sup_{W \cap P_x} |f_k| - \sup_{\partial(W \cap P_x)} |f_k|}{\text{diam}(W \cap P_x)} \right|^d dx &\leq \int_M \int_{W \cap P_x} |\det D^2(f_k|_{P_x})| dx \\ &\leq \nu_{k,d}(W). \end{aligned}$$

By dominated convergence,

$$\begin{aligned} \int_M \left| \frac{\sup_{W \cap P_x} |f| - \sup_{\partial(W \cap P_x)} |f|}{\text{diam}(W \cap P_x)} \right|^d dx &= \lim_{k \rightarrow \infty} \int_M \left| \frac{\sup_{W \cap P_x} |f_k| - \sup_{\partial(W \cap P_x)} |f_k|}{\text{diam}(W \cap P_x)} \right|^d dx \\ &\leq \limsup_{k \rightarrow \infty} \nu_{k,d}(W) \\ &\leq \nu_d(V). \end{aligned}$$

Taking $W = W_1, W_2, \dots$ to be an exhaustion of V , dominated convergence implies that the left hand side converges to the corresponding expression for V . $\square$

3. AN EXTENSION OF A SPECIAL CASE OF A THEOREM OF CALDERÓN AND ZYGMUND

We say that f **admits an approximate second derivative** at x if (1) holds with f replaced by the restriction $f|_E$, where E is some measurable set with density 1 at x . We say that f **admits a k th derivative in the L^1 sense** at x if there exists a polynomial Q_x of degree k such that

$$(9) \quad r^{-n} \int_{B_r(x)} |f(y) - Q_x(y)| dy = o(r^k)$$

as $r \downarrow 0$. An easily proved equivalent formulation for first derivatives may be stated as follows.

Lemma 3.1. *The function g is differentiable at 0 in the L^1 sense iff the one-parameter family of functions $g_r(x) := rg(r^{-1}x)$ converges in L^1_{loc} to a linear function.*

The following is also obvious.

Lemma 3.2. *If f admits a second derivative in the L^1 sense at x , then f admits an approximate second derivative at x .*

The next proposition generalizes a result of Calderón-Zygmund [3]. The original result of [3] (or rather the very special case of it that we have in mind) states that a function with distributional second derivatives in L^1_{loc} admits a second derivative in the L^1 sense a.e. By a straightforward adaptation of the argument of [3] we prove that this conclusion is true if the distributional second derivatives are only locally finite signed measures. We recall that the space $\text{BV}_{\text{loc}}(U)$ of functions of locally bounded variation consists of all locally integrable functions whose distributional gradients are (vector) measures (cf. [8]).

Proposition 3.3. *If the distributional gradient of $f \in L^1(U)$ lies in $\text{BV}_{\text{loc}}(U)$, then f admits a second derivative in the L^1 sense a.e. in U .*

Lemma 3.4. *If $g \in \text{BV}_{\text{loc}}(U)$ then g is differentiable in the L^1 sense at a.e. $x \in U$.*

Proof. By [4], Thm. 3.22, in the Lebesgue decomposition of the distributional gradient ∇g into its singular and absolutely continuous parts with respect to Lebesgue measure, the singular part has density zero at a.e. $x \in U$. Let $x = 0$ be such a point, and assume further that 0 is a Lebesgue point for g and for the absolutely continuous part of ∇g . We may assume without loss of generality that $g(0) = 0$.

By the BV compactness theorem (cf. [8], Theorem 1.9), for some sequence $r_i \downarrow 0$ the dilates $G_i(x) := r_i g(r_i^{-1}x)$ converge in $L^1_{\text{loc}}(\mathbb{R}^n)$ to some function $\lambda \in L^1_{\text{loc}}(\mathbb{R}^n)$. Furthermore the distributional gradients ∇G_i converge weakly to the constant $\nabla g(0)$. Therefore λ is the linear function $\lambda(x) := \sum_{i=1}^n \frac{\partial g}{\partial x_i}(0)x_i$.

Since this conclusion is independent of the sequence r_i , in fact $rg(r^{-1}x) \rightarrow \lambda(x)$ in L^1_{loc} , which by Lemma 3.1 is equivalent to the stated conclusion. $\square$

Proof of Prop. 3.3. By Lemma 3.4 it is enough to show that there exists an L^1 quadratic Taylor approximation for f at 0 provided ∇f is differentiable in the L^1 sense at 0. We may assume also that $f(0) = \nabla f(0) = D^2 f(0) = 0$, where $D^2 f(0)$ is the L^1 derivative of ∇f at 0. Put

$$G(\rho) := \int_{B(0,\rho)} \frac{|\nabla f(x)|}{|x|^{n-1}} dx, \quad F(\rho) := \int_{B(0,\rho)} |\nabla f(x)| dx.$$

Then F, G are both absolutely continuous on $[0, \infty)$, with $G'(\rho) = \frac{F'(\rho)}{\rho^{n-1}}$, and $F(\rho) = o(\rho^{n+1})$ by Lemma 3.4. Integrating by parts, it follows that $G(\rho) = o(\rho^2)$ as $\rho \downarrow 0$. On the other hand,

$$\begin{aligned} \int_{B(0,\rho)} |f| &\leq C \int_0^\rho r^{n-1} dr \int_{S^{n-1}} dv \int_0^r |Df(sv)| ds \\ &\leq C \int_0^\rho r^{n-1} dr \int_{S^{n-1}} dv \int_0^\rho |Df(sv)| ds \\ &= C \rho^n \int_{S^{n-1}} dv \int_0^\rho |Df(sv)| ds \\ &= C \rho^n G(\rho) = o(\rho^{n+2}), \end{aligned}$$

which gives the result. $\square$

4. PROOF OF THEOREM 1.1

To prove Theorem 1.1 we use induction on the dimension n . For $n = 1$, the result follows at once from Alexandrov's theorem and the parenthetical remark following the statement of the Theorem 1.1. Alternatively, it is also easy to prove by direct integration that f is twice differentiable at every point at which f' is differentiable in the L^1 sense. Supposing this to be true at 0, we may assume that the L^1 second derivative at 0 is 0, so the relation (9) yields

$$f(x) = \int_0^x f'(s) ds = o(x^2),$$

i.e. (1) holds at 0 with $Q_0 = 0$.

4.1. Setting up the inductive step. Put H for the space of all hyperplanes in $\mathbb{R}^n$, not necessarily passing through the origin, and $I := \{(x, P) \in \mathbb{R}^n \times H : x \in P\} \simeq \mathbb{R}^n \times \mathbb{RP}^{n-1}$. Thus I admits the double fibration

$$(10) \quad \mathbb{R}^n \xleftarrow{p} I \xrightarrow{q} H.$$

Furthermore I admits a measure μ , invariant with respect to euclidean motions, expressible in local coordinates for either bundle structure as the product of invariant measures on the base and the fiber.

Lemma 4.1. *Put E for the set of pairs $(x, P) \in I$ such that $f|P$ is twice differentiable at x . Then E is a Borel subset of I .*

Proof. For $i, j, k \in \mathbb{N}$, define

$E_{ijk} := \{(x, P) \in I : \text{there exists a quadratic polynomial } Q, \|Q\| \leq k, \text{ such that } |f(x') - Q(x')| \leq \frac{1}{i}|x' - x|^2 \text{ whenever } x' \in P \text{ and } |x' - x| < \frac{1}{j}\}.$

We claim first that each E_{ijk} is a closed subset of I . Suppose $E_{ijk} \ni (x_m, P_m) \rightarrow (x_0, P_0)$ and $x'_0 \in P_0, |x'_0 - x_0| < \frac{1}{j}$. Let Q_m be a quadratic polynomial as in the definition of E_{ijk} for (x_m, P_m) . We may assume that the sequence of the Q_m converges to a quadratic polynomial Q_0 with $\|Q_0\| \leq k$. Clearly there exist points $x'_m \in P_m$ with $|x'_m - x_m| < \frac{1}{j}$ and $x'_m \rightarrow x'_0$ (e.g., for large m we may take x'_m to be the orthogonal projection of x'_0 to P_m) so

$$|f(x'_0) - Q_0(x'_0)| = \lim |f(x'_m) - Q_m(x'_m)| \leq \frac{1}{i} \lim |x'_m - x_m|^2 = \frac{1}{i} |x'_0 - x_0|^2.$$

Thus $(x_0, P_0) \in E_{ijk}$.

Next we claim that

$$E = \bigcup_k \bigcap_i \bigcap_j E_{ijk} =: F,$$

which is sufficient to establish the desired conclusion.

That $E \subset F$ is obvious. To prove that $F \subset E$, suppose $(x, P) \in F$. Since $\|Q(\cdot - x)\| \leq (1 + 2|x| + |x|^2) \|Q\|$, we may suppose that $x = 0$. Fix k , and suppose that $(0, P) \in \bigcap_i \bigcup_j E_{ijk}$. Then for each i there is j_i such that $(0, P) \in E_{ij_i k}$; let Q_i be the quadratic polynomial in the corresponding defining condition. Clearly the restrictions $Q_i|_P$ all agree to first order at 0, and $|(Q_i - Q_{i'})(v)| \leq \frac{2|v|^2}{\min(i, i')}$ for $v \in P$. Altering the Q_i off of P if necessary, we may assume that this is true for all $v \in \mathbb{R}^n$. Thus the Q_i converge to a quadratic polynomial Q_0 , and for $x' \in P, |x'| < \frac{1}{j_i}$,

$$\begin{aligned} |f(x') - Q_0(x')| &\leq |f(x') - Q_i(x')| + |Q_i(x') - Q_0(x')| \\ &\leq \frac{1}{i}|x'|^2 + \frac{2}{i}|x'|^2 = \frac{3}{i}|x'|^2 \end{aligned}$$

It follows that $f|P$ is twice differentiable at 0, with second order Taylor expansion Q_0 , which establishes the claim. $\square$

Assume the conclusion of Theorem 1.1 is true for dimension $n - 1$, and let $x_0 \in \mathbb{R}^n$ be a point at which f is approximately twice differentiable—this is true for a.e. $x_0 \in U$ by Lemma 3.2 and Prop. 3.3. We may assume further that $f|P$ is twice differentiable at x_0 for almost every hyperplane P passing through x_0 : it is clear that for a.e. P , the hypotheses of Theorem 1.1 hold for some subsequence $f_{k'}|P \rightarrow f|P$, hence $f|P$ is twice differentiable a.e. in P by induction. Thus by Lemma 4.1 and Fubini's theorem, the set $\{(x, P) : f|P \text{ is twice differentiable at } x\}$ has full measure in I . Applying Fubini's theorem again, we find that for a.e. $x \in \mathbb{R}^n$, the set $\{P : f|P \text{ is twice differentiable at } x\}$ has full measure in the space of hyperplanes passing through x . We will show that if x_0 is such a point, and if f does *not* admit a second order Taylor expansion at x_0 , then at least one of the

absolute Hessian determinant measures $\nu_0, \dots, \nu_n$ has upper density $\Theta^*(\nu_i, x_0) = \infty$ at x_0 . This will conclude the proof.

We may assume that $x_0 = 0$. We may assume also that the second order approximate Taylor expansion Q of f at 0 is identically zero: otherwise, in light of Lemma 2.1, we may replace f by $f - Q$. With these assumptions we have

Lemma 4.2. *For a.e. hyperplane P passing through 0, the second order Taylor expansion Q^P of $f|_P$ at 0 is zero.*

Proof. Let $E \subset U$ be the subset of density 1 at 0 in the definition of the approximate second derivative at the beginning of section 3. Since f is continuous, we may assume that E is closed.

We claim that a.e. line l through 0 meets E at points lying arbitrarily close to 0, i.e. $l \cap B(0, r) \cap E - \{0\} \neq \emptyset$ for all $r > 0$. In other words, putting $A_r := \{l : l \cap B(0, r) \cap E - \{0\} \neq \emptyset\}$ for $r > 0$ and $A_0 := \bigcap_{r>0} A_r$, where $A_r \downarrow A_0$ as $r \downarrow 0$, we claim that A_0 has full measure in the space of lines through 0. If this were not the case then there would exist some $r_0 > 0$ such that A_{r_0} does not have full measure, from which it follows that for $r < r_0$

$$\frac{\text{vol}(B(0, r) \cap E)}{\omega_n r^n} \leq \frac{\text{vol}\left(B(0, r_0) \cap \bigcup_{l \in A_{r_0}} l\right)}{\omega_n r_0^n} < 1,$$

which is a contradiction.

Now, if $l \in A_0$ then $\liminf_{v \rightarrow 0, v \in l} \frac{|f(v)|}{|v|^2} = 0$. Hence if $v \in l \subset P$, and $f|_P$ is twice differentiable at 0, then $Q^P(v) = 0$. But, by basic integral geometry, a.e. P through 0 has the property that a.e. line $l \subset P$ through 0 belongs to A_0 . Therefore $Q^P \equiv 0$ on a dense subset of P , and therefore $Q^P \equiv 0$ by continuity. $\square$

4.2. The main lemma. For $\epsilon > 0$ we consider the open sets

$$(11) \quad V_\epsilon := \{x : |f(x)| > \epsilon|x|^2\}$$

The approximate twice differentiability hypothesis means that each V_ϵ has density 0 at 0. The statement that f is twice differentiable at 0, with zero 2nd order Taylor expansion there, is equivalent to the statement that for each $\epsilon > 0$ the set $V_\epsilon \cap B(x, r) = \emptyset$ for all sufficiently small $r > 0$.

Lemma 4.3. *Suppose the upper densities $\Theta^*(\nu_i, 0) < \infty, i = 0, \dots, n-1$. Then for fixed $\epsilon > 0$, the supremum of the diameters of the components of $B(0, r) \cap V_\epsilon$ is $o(r)$ as $r \downarrow 0$.*

Proof of Lemma 4.3. Put I_0 for the space of hyperplanes through 0. By Lemma 4.2, for a.e. $P \in I_0$ there exists $r = r(P) > 0$ such that

$$(12) \quad P \cap V_\epsilon \cap B(0, r) = \emptyset.$$

For $r > 0$, put

$$(13) \quad \Pi_r := \{P : P \in I_0, P \cap V_\epsilon \cap B(0, r) \neq \emptyset\}.$$

Thus the family of subsets $\Pi_r \subset I_0$ is decreasing as $r \downarrow 0$, with

$$(14) \quad \lim_{r \downarrow 0} \mu_0(\Pi_r) = \mu_0\left(\bigcap_{r>0} \Pi_r\right) = 0,$$

where μ_0 is the invariant probability measure on I_0 .

Now consider the spherical open sets

$$(15) \quad W_{r,\epsilon} := \{v \in S^{n-1} : \text{there exists } r' \in (0, r) \text{ such that } r'v \in V_\epsilon\},$$

i.e. $W_{r,\epsilon}$ is the spherical projection of $B(0, r) \cap V_\epsilon$. Thus $P \in \Pi_r$ iff $P \cap W_{r,\epsilon} \neq \emptyset$. It now follows from (14) and elementary integral geometry (cf. [12]) that the supremum of the spherical diameters of the components of $W_{r,\epsilon}$ converges to 0 as $r \downarrow 0$ (ϵ fixed)

To complete the proof of the lemma, it is now enough to show that the “radial diameter” $\sup_C |x| - \inf_C |x| = o(r)$ for any component C of $B(0, r) \cap V_\epsilon$.

Let $\delta < \frac{\pi}{6}$ be given, and take $r_0 > 0$ small enough that

- (1) each component of $W_{\frac{2r_0}{\sqrt{3}}, \frac{\epsilon}{2}} \subset S^{n-1}$ is included in a spherical ball of radius δ , and
- (2) $\frac{\nu_{n-1}(B(0, 2r_0/\sqrt{3}))}{\omega_n(2r_0/\sqrt{3})^n} < \Theta^*(\nu_{n-1}, 0) + 1$.




FIGURE 1. Proof of Lemma 4.3

Let C be any component of $V_\epsilon \cap B(0, r_0)$, and C' the component of $V_{\epsilon/2} \cap B(0, 2r_0/\sqrt{3})$ that includes C . Assumption (1) above guarantees that there is a spherical ball of radius δ that includes the spherical projection of C' . Let v_0 be its center and put for $t \in \mathbb{R}$

$$P_t := \{x : \langle x, v_0 \rangle = t\}.$$

Finally, let $(a, b) := \{t \in (0, r_0) : P_t \cap C \neq \emptyset\}$. Thus for $t \in (a, b)$

$$(16) \quad \sup_{C' \cap P_t} |f| = \sup_{C \cap P_t} |f| > \epsilon t^2.$$

By assumption (1),

$$(17) \quad \sup_C |x| - \inf_C |x| < (b - a) \sec \delta < (b - a) \frac{2}{\sqrt{3}}$$

and, if $x \in C' \cap P_t$,

$$(18) \quad t \leq |x| < t \sec \delta < \frac{2t}{\sqrt{3}}.$$

In particular, if $P_t \cap B(0, r_0) \neq \emptyset$ (i.e. if $t < r_0$) then $C' \cap P_t \subset \subset B(0, 2r_0/\sqrt{3})$, and in particular

$$|f(x)| = \frac{\epsilon |x|^2}{2} < \frac{2\epsilon t^2}{3}$$

for $x \in \partial C' \cap P_t$. Thus we may apply Corollary 2.3 to obtain

$$\begin{aligned} (\Theta^*(\nu_{n-1}, 0) + 1) \omega_n \left(\frac{2r_0}{\sqrt{3}} \right)^n &> \nu_{n-1} \left(B(0, \frac{2r_0}{\sqrt{3}}) \right) \\ &\geq \int_a^b \left| \frac{\sup_{C' \cap P_t} f - \sup_{\partial C' \cap P_t} f}{\text{diam}(C' \cap P_t)} \right|^{n-1} dt \\ &\geq \int_a^b \left(\frac{\epsilon t^2 - \frac{2}{3}\epsilon t^2}{2t \tan \delta} \right)^{n-1} dt \\ &= \left(\frac{\epsilon}{6} \right)^n \cot^n \delta \frac{(b^n - a^n)}{n} \\ &\geq \left(\frac{\epsilon}{6} \right)^n \cot^n \delta \frac{(b - a)^n}{n}. \end{aligned}$$

Therefore, by (17),

$$\sup_C |x| - \inf_C |x| \leq \frac{2}{\sqrt{3}}(b - a) < \frac{C}{\epsilon} (\Theta^*(\nu_{n-1}, 0) + 1)^{\frac{1}{n}} r_0 \tan \delta.$$

Since $\delta > 0$ can be taken arbitrarily small (and ϵ is fixed), this completes the proof of the lemma. $\square$

4.3. Completion of the proof Theorem 1.1. Suppose $x_1, x_2, \dots \in V_\epsilon$ with $x_i \rightarrow 0$. Let C_i be the component of $V_\epsilon \cap B(0, 2|x_i|)$ containing x_i , and C'_i the component of $V_{\frac{\epsilon}{2}} \cap B(0, 2|x_i|)$ that includes C_i . By Lemma 4.3, the diameter of C'_i is $o(|x_i|)$ as $i \rightarrow \infty$, hence $C'_i \subset \subset B(0, 2|x_i|)$ for i sufficiently large. Applying Lemma 2.2,

$$\begin{aligned} \frac{\nu_n(B(0, 2|x_i|))}{|x_i|^n} &\geq \frac{\nu_n(C'_i)}{|x_i|^n} \\ &\geq \left| \frac{\sup_{C'_i} |f| - \sup_{\partial C'_i} |f|}{|x_i| o(|x_i|)} \right|^n \\ &\geq \left(\frac{\epsilon}{2} \right)^n \left(\frac{|x_i|}{o(|x_i|)} \right)^n \\ &= \frac{1}{o(1)} \rightarrow \infty \end{aligned}$$

as $i \rightarrow \infty$ since ϵ is fixed. Therefore the upper density of ν_n at 0 is infinite.

4.4. A concluding remark. In fact our proof yields a slightly more general statement than Theorem 1.1: besides continuity, the only properties of f that we have used are the existence of the absolute Hessian determinant measures, the transformation law of Lemma 2.1, and their role in Lemmas 2.2 and 2.3.

REFERENCES

- [1] Alberti, G., Ambrosio, L. A geometrical approach to monotone functions in $\mathbb{R}^n$. *Math. Z.* 230 (1999), pp. 259–316.
- [2] Aleksandrov, A.D. Almost everywhere existence of the second differential of a convex function and some properties of convex surfaces connected with it. *Leningrad Univ. Ann. (Math. ser.)* **6** (1939), 335 (in Russian).
- [3] Calderon, A.P. and Zygmund, A. Local properties of elliptic partial differential equations. *Studia Math.* **20** (1961), 171–225
- [4] Folland, G.B. *Real analysis*. Wiley, New York, 1999.
- [5] Fu, J. H. G.: Monge-Ampère functions I. *Indiana Univ. Math. J.* **38** (1989), 745–771
- [6] Fu, J. H. G.: Monge-Ampère functions II. *Indiana Univ. Math. J.* **38** (1989), 773–789
- [7] Gilbarg, D. and Trudinger, N. *Elliptic Partial Differential Equations of Second Order*. Springer, New York, 1983
- [8] Giusti, E. *Minimal Surfaces and Functions of Bounded Variation*. Birkhäuser, Boston, 1984
- [9] Hutchinson, J.E. and Meier, M. A remark on the nonuniqueness of tangent cones. *Proc. Amer. Math. Soc.* 97 (1986), pp. 184–185.
- [10] Jerrard, R.L.: Some remarks on Monge-Ampère functions. *Singularities in PDE and the Calculus of Variations*, Stanley Alama, Lia Bronsard, and Peter J. Sternberg, eds. - AMS — CRM, 2008
- [11] Jerrard, R.L.: Some rigidity results related to Monge-Ampère functions. To appear in *Canadian Journal of Mathematics*
- [12] Santaló, L. *Integral geometry and geometric probability*, 2nd ed. Cambridge University Press, Cambridge 2004

E-mail address: fu@math.uga.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS, GA 30602, USA